PHOTOMETRIC OBSERVATION AND ANALYSIS OF THE CLOSE BINARY SYSTEM BO PEGASI

ATSUMA YAMASAKI*†

Department of Earth Science and Astronomy, University of Tokyo, Meguro-ku, Tokyo 153, Japan

AKIRA OKAZAKI*†

Department of Mathematics, Tsuda College, Kodaira, Tokyo 187, Japan Received 1986 July 24, revised 1986 September 11

ABSTRACT

Photoelectric BV light curves of the close binary system BO Peg (P=0.580) are obtained and analyzed with a synthesis method to determine the photometric elements. It is found that the components of BO Peg are not in contact with each other but almost fill their respective Roche lobes. Two spectrograms of the system (A7 IV–V) have also been obtained, and a brief discussion on physical quantities of the components is made.

Key words: close binary-photometry-light-curve analysis

I. Introduction

The eclipsing binary BO Pegasi was first discovered by Hoffmeister (1935). Jensch (1935) presented the photographic light curve and determined the orbital period to be 0.458044. The spectral type was given as A4 in the General Catalogue of Variable Stars (Kukarkin et al. 1971). No photoelectric light curves have been published so far.

In this study, we present photoelectric *BV* light curves, analyze the light curves, determine a new spectral type from our spectrograms, and discuss briefly the physical properties of BO Peg.

II. Photoelectric Observations

We made photoelectric observations of BO Peg with the 0.6-m reflector of the University of Hawaii at Mauna Kea Observatory on six nights in September 1984. We employed the Tinsley-Goguen photometer with a refrigerated photomultiplier RCA C 31034A(GaAs) and Apple Data System (Goguen 1984). Most of the observations were carried out in *B* and *V*, and a few observations in *U* were also made. The filters used are very similar to those of Johnson's *UBV* standard system.

Two nearby field stars were respectively chosen as the comparison star and the check star. The observations showed no appreciable variations in differential magnitudes between the comparison star and the check star.

Table I gives the positions, magnitudes, and color indices of the comparison star and the check star as well as BO Peg at maximum.

We secured the time of the primary minimum, HJD2445969.8604 with (O-C) = -0.0399 from the ephemeris Min. I = HID2427655.460 + 0.5804342 E(Whitney 1957). Table II lists available observed times of minima of BO Peg, including the present one. From these 28 minima, we derived the ephemeris Min. I = HJD2445969.8709 + 0.58043326 E, by using the leastsquares method with weights 1, 3, and 10 for visual, photographic, and photoelectric observations, respectively. The (O-C) values calculated from the latter ephemeris are given in the third column of Table II and are plotted in Figure 1. It is found that neither the ephemeris of Whitney (1957) nor the revised one satisfies the present light variations of BO Peg: the two opposite trends in the (O-C) diagram around E=-20,000 and 0 suggest that period changes may have occurred. We need more photoelectric observations.

Differential magnitudes ΔV , ΔB , and ΔU , in the sense

TABLE 1
Positions, Magnitudes, and Color Indices of BO Peg, Comparison Star, and Check Star

Star	RA(1950)	Dec(1950)	V B - V		U - B	
BO Peg	21 ^h 28 ^m 54 ^s	+11°43.6	11.90*	+0.36*	+0.10*	
Comparison	21 29 26	+11 41.9	9.73	+0.51	+0.07	
Check	21 29 36	+11 39.9	9.87	+0.50	+0.02	

^{*} At maximum light.

^{*}Guest Observer, Mauna Kea Observatory, Institute for Astronomy, University of Hawaii.

 $[\]dagger$ Observations were partially made at Okayama Astrophysical Observatory.

TABLE []
Observed Times of Primary Minima and Their (O-C) Values

HJD2400000+	E	o-c*	Method	Source**
00000 501	0.4.0.0.0	0 ^d 000		
26223.531	-34020		pg	1
26244.392	-33984	-0.035	pg	1
26333.238	-33831	+0.005	pg	1
27655.450	-31553	-0.010	vis	1
27658.362	-31548	0.000	vis	1
27666.500	-31534	+0.012	vis	1
27684.490	-31503	+0.008	vis	1
27685.630	-31501	-0.013	vis	1
27688.547	-31496	+0.002	vis	1
27691.450	-31491	+0.003	vis	1,
27699.580	-31477	+0.007	vis	1
27745.426	-31398	-0.001	vis	1
32005.793	-24058	-0.015	pg	
32799.842	-22690	+0.002	pg	2 2 2 2 2 2 2
33584.599	-21338	+0.013	pg	$\overline{2}$
33981.612	-20654	+0.010	pg	2
34245.705	-20199	+0.006	pg	2
34627.642	-19541	+0.017	pg	2
35002.600	-18895	+0.016	pg	2
44194.337	-3059	+0.011	vis	3
44201.296	-3047	+0.005	vis	3
44458.426	-2604	+0.003	vis	4
44458.439	-2604	+0.016	vis	4
44490.354	-2549	+0.008	vis	5
44490.355	-2549	+0.009	vis	5
45612.323	-616	-0.001	vis	6
45969.8604	0	-0.0105	pe	This study
46298.388	+566	-0.008	vis	7

* The O-C values are calculated with the ephemeris: Min I = HJD2445969.8709 + 0.58043326 E.

** Sources. 1: Jensch 1935, 2: Whitney 1957.
3: BBSAG (Bedeckungsveränderlichen Beobachter der Schweizerischen Astronomischen Gesellschaft) Bull., No.45 (1979), 4: BBSAG Bull., No.49 (1980), 5: BBSAG Bull., No.50 (1980), 6: BBSAG Bull., No.69 (1983), 7: BBSAG Bull., No.78 (1985).

variable minus comparison, are given in Appendix Table A1, where phases are calculated using the ephemeris

Min. I = HID
$$2445969.8604 + 0.58043288 E$$
,

which was derived by adopting the time of the primary minimum obtained in this study as the epoch. The light variations in V and B are also shown in Figure 2. It is obvious that this system suffers large proximity effects.

III. Light-Curve Analysis

We analyzed the *BV* light curves using the light-curve synthesis method developed by one of us (A.Y.). In this method, the photospheric configurations of the components are approximated by the Roche equipotential surfaces (e.g., Yamasaki, Okazaki, and Kitamura 1983).

In the present analysis, the surfaces of the primary and secondary components were divided into 994 and 582 elements, respectively. The envelope of the primary component was assumed to be radiative and that of the secondary component to be convective. We adopted the following values throughout the analysis:




Fig. 1-(O-C) values of times of minima of BO Peg. Dots, open circles, and dot-open circle denote visual, photographic, and photoelectric observations, respectively.




FIG. 2-Light variations of BO Peg in V and B. Theoretical curves are shown with solid curves.

,	Primary	Secondary
Limb-darkening coefficient:	$u_{\rm IV}=0.56$	$u_{2V}=0.60$
	$u_{1B}=0.68$	$u_{2B}=0.74$
Gravity-darkening exponent:	$\beta_1 = 0.25$	$\beta_2 = 0.08$
Albedo:	$A_1 = 1.0$	$A_0 = 0.5$.

where the linear limb-darkening coefficients of the primary component are for an \sim F0-type star and those of the secondary component are for an \sim G0-type star (Carbon and Gingerich 1969).

The system parameters have been obtained by minimizing $(O-C)^2 = \sum_i (l_{\rm obs} - l_{\rm th})_i^2$, where the summation is over all observations. Table III gives the final solution of the system parameters for BO Peg. Theoretical light curves computed with these parameters are shown for V and B in Figure 2 with solid curves.

Figure 3 represents the configurations of the components of BO Peg relative to their critical Roche lobes. It is shown that the components are not in contact with each

BO PEGASI 1327

TABLE III

System Parameters of BO Peg

Mass ratio: q		0.55 ± 0.05
Inclination: i		79°1 ±1°0
Fractional radius of the primary component: r ₁ (Roche,	side substellar back pole side	0.430 ±0.010 0.545 0.458 0.406 0.431)
Fractional radius of the secondary component: r ₂ (Roche,	side substellar back pole side	0.310 ±0.010 0.373 0.337 0.298 0.321)
Fractional luminosity of the primary component: L_1	V B	0.914 ±0.020 0.919 ±0.020
Fractional luminosity of the secondary component: L_2	V B	0.086 ±0.020 0.081 ±0.020

other, although they almost fill their critical Roche lobes—the primary component fills its critical Roche lobe and the secondary component is very close to its critical lobe.

IV. Spectroscopic Observations

We secured two spectrograms of BO Peg with a dispersion of 32 Å mm⁻¹ with the coudé image-tube spectrograph of the 1.9-m reflector at Okayama Astrophysical Observatory in November 1980. These spectrograms were taken outside eclipses on HJD2444545.008 (0.190) and HJD2444547.067 (0.737). No emission features were detected in these spectrograms. The spectral type of BO Peg outside eclipses was determined to be A7 IV–V or slightly later.

V. Discussion

To deduce reliable absolute dimensions of BO Peg, radial-velocity observations are essential; however, it is not practical to make discussions with only two spectrograms. Therefore, in the following discussions, we assume the mass of the primary component $\mathfrak{M}_1=1.9~\mathfrak{M}_{\odot}$ corresponding to the spectral type of A7 IV–V (Allen 1973). From the system parameters given in Table III, we obtain:

Both components of BO Peg are found to have radii




FIG. 3—Configurations of the components of BO Peg (solid line) relative to their critical Roche lobes (dotted line). P and S represent the centers of the primary and the secondary components, respectively.

slightly larger than those of corresponding main-sequence stars of the same masses. The secondary component is considered to be an early-G-type star from its mass.

According to the light-curve analysis in Section III, the components are not in contact, although they are very close to each other. From the fractional luminosities and radii in Table III, the brightness ratio J_1/J_2 was derived to be 5.5–5.9. The large brightness ratio also suggests that the system of BO Peg does not have a common envelope.

In conclusion, BO Peg is a new member of short-period noncontact binary systems (Yamasaki *et al.* 1983; Yamasaki, Okazaki, and Kitamura 1984), or almost contact systems, in which interacting-binary evolution off the main sequence has just taken place.

We would like to thank Professor D. N. B. Hall, Director, Professor J. P. Henry, and the staff of the University of Hawaii at Mauna Kea Observatory. We are grateful to Dr. J. Goguen for the photometric system. We also thank the staff of the Okayama Astrophysical Observatory. This work was supported by the Grant-in-Aid for Overseas Scientific Survey (59041022, 60043023) of the Ministry of Education, Science, and Culture of Japan.

REFERENCES

Allen, C. W. 1973, Astrophysical Quantities (3d ed.; London: Athlone Press), p. 209.

Carbon, D. F., and Gingerich, O. 1969, in Theory and Observation of Normal Stellar Atmospheres, ed. O. Gingerich (Cambridge, MA: MIT Press), p. 377.

Goguen, J. 1984, unpublished.

Hoffmeister, C. 1935, Astr. Nach., 255, 401.

Jensch, A. 1935, Astr. Nach., 255, 417.

Kukarkin, B. V., et al. 1971, First Supplement to the Third Edition of the General Catalogue of Variable Stars (Moscow: Academy of Sciences of the USSR).

Whitney, B. S. 1957, A.J., 62, 371.

Yamasaki, A., Okazaki, A., and Kitamura, M. 1983, Pub. Astr. Soc. Japan, 35, 131.

_____. 1984, Pub. Astr. Soc. Japan, 36, 175.

YAMASAKI AND OKAZAKI

TABLE A1
Photoelectric Observations of BO Peg

HJD 2445000+	Phase	Δ۷	HJD 2445000+	Phase	ΔV	HJD 2445000+	Phase	ΔВ	HJD 2445000+	Phase	ΔВ
958.8432	0.0190	2.891	971.0034	0.9692	2.759	958.8901	0.0998	2.262	971.0336	0.0213	2.788
958.8630	0.0530	2.706	971.0091	0.9790	2.846	958.9001	0.1170	2.196			
958.8720	0.0687	2.572	971.0166	0.9920	2.907 2.916	958.9080	0.1307	2.154	971.7246	0.2117	2.025
958.8805 958.8891	0.0833	2.480 2.407	971.0198 971.0225	0.9975	2.918	958.9163 958.9248	0.1449	2.125 2.113	971.7322 971.7492	0.2249	2.038
958.8991	0.1152	2.341	971.0267	0.0094	2.927	958.9332	0.1741	2.073	971.7561	0.2660	2.033
958.9070	0.1289	2.305	971.0326	0.0196	2.895	958.9407	0.1870	2.078	971.7607	0.2739	2.025
958.9152	0.1431	2.273				958.9484	0.2002	2.065	971.7648	0.2810	2.024
958.9238	0.1579	2.271	971.7236	0.2100	2.181	958.9563	0.2139	2.048	971.7695	0.2891	2.033
958.9322	0.1724	2.222	971.7312	0.2232	2.202	958.9642	0.2274	2.009	971.7736	0.2962	2.040
958.9397	0.1852	2.230	971.7483	0.2525	2.179	958.9725	0.2418	2.019	971.7776	0.3031	2.051
958.9474 958.9553	0.1984	2.210 2.198	971.7552 971.7600	0.2645	2.182 2.183	958.9820 958.9897	0.2582	2.004 2.002	971.7816 971.7858	0.3099	2.047
958.9632	0.2258	2.173	971.7640	0.2797	2.171	958.9971	0.2841	2.018	971.7900	0.3171	2.069
958.9715	0.2400	2.158	971.7687	0.2878	2.192	959.0048	0.2974	2.028	971.7944	0.3321	2.073
958.9811	0.2565	2.152	971.7730	0.2951	2.194	959.0323	0.3448	2.104	971.7983	0.3388	2.083
958.9886	0.2696	2.167	971.7769	0.3019	2.202				971.8026	0.3461	2.088
958.9960	0.2822	2.149	971.7809	0.3087	2.192	967.7607	0.3826	2.134	971.8067	0.3531	2.098
959.0038	0.2957	2.177	971.7850	0.3158	2.218	967.7681	0.3953	2.161	971.8119	0.3622	2.120
959.0289	0.3390	2.248	971.7892	0.3230	2.211	967.7768 967.7836	0.4102	2.172 2.238	971.8170	0.3710	2.122
967.7597	0.3809	2.281	971.7937 971.7976	0.3375	2.217 2.241	707.7030	0.4217	2.230	971.8209 971.8249	0.3776 0.3845	2.136
967.7663	0.3921	2.309	971.8018	0.3448	2.234	969.7439	0.7993	2.041	971.8289	0.3915	2.154
967.7758	0.4085	2.313	971.8059	0.3519	2.256	969.7495	0.8089	2.042	971.8330	0.3985	2.170
967.7825	0.4201	2.370	971.8117	0.3617	2.258	969.7549	0.8183	2.053	971.8371	0.4055	2.191
			971.8164	0.3698	2.272	969.7603	0.8276	2.069	971.8410	0.4122	2.200
969.7429	0.7976	2.195	971.8202	0.3764	2 280	969.7657	0.8369	2.084	971.9228	0.5532	2.297
969.7485	0.8072	2.201	971.8241	0.3832	2.282	969.7711	0.8461	2.100	971.9278	0.5619	2.287
969.7539	0.8166	2.212	971.8282	0.3902	2.305	969.7764	0.8554	2.109	971.9321	0.5693	2.271
969.7593 969.7648	0.8258 0.8352	2.238 2.242	971.8322 971.8364	0.3971	2.323	969.7817 969.7874	0.8644 0.8742	2.119 2.122	971.9363	0.5764	2.252
969.7701	0.8444	2.251	971.8304	0.4115	2.353	969.8450	0.9735	2.684	971.9408 971.9452	0.5841	2.242
969.7755	0.8537	2.260	971.9222	0.5521	2.450	969.8502	0.9824	2.746	971.9496	0.5994	2.212
969.7807	0.8627	2.272	971.9272	0.5608	2.441	969.8557	0.9919	2.794			
969.7864	0.8725	2.292	971.9313	0.5679	2.427	969.8615	0.0018	2.806	972.8651	0.1767	2.072
969.8441	0.9718	2.784	971.9356	0.5753	2.408	969.8666	0.0108	2.800	972.8695	0.1841	2.068
969.8493	0.9808	2.859	971.9400	0.5828	2.369	969.8707	0.0178	2.781	972.8738	0.1916	2.055
969.8547 969.8604	0.9902	2.896 2.919	971.9446 971.9489	0.5907 0.5982	2.376 2.372	969.8762 969.8817	0.0272 0.0368	2.735 2.667	972.8782 972.8827	0.1992 0.2070	2.053
969.8657	0.0091	2.914	7/1.7407	0.3702	2.3/2	969.8870	0.0459	2.595	972.8871	0.2146	2.035
969.8698	0.0161	2.896	972.8644	0.1754	2.240	969.8949	0.0594	2.500	972.8981	0.2335	2.012
969.8752	0.0255	2.855	972.8687	0.1828	2.224	969.9004	0.0689	2.413	972.9051	0.2456	2.027
969.8808	0.0351	2.802	972.8730	0.1903	2.208	969.9487	0.1521	2.118	972.9117	0.2569	2.009
969.8861	0.0442	2.740	972.8774	0.1979	2.208	969.9545	0.1622	2.094	972.9720	0.3608	2.107
969.8939	0.0578	2.646	972.8820	0.2058	2.196	969.9602	0.1719	2.080	972.9770	0.3694	2.121
969.8994 969.9477	0.0672 0.1504	2.557 2.271	972.8864 972.8973	0.2133	2.187 2.157	969.9657 969.9718	0.1815 0.1919	2.060 2.049	972.9813 972.9858	0.3768 0.3846	2.143
969.9536	0.1606	2.250	972.9043	0.2442	2.178	969.9800	0.2060	2.036	972.9904	0.3925	2.168
969.9592	0.1702	2.237	972.9110	0.2557	2.174	969.9856	0.2156	2.028	972.9962	0.4025	2.196
969.9647	0.1798	2.224	972.9713	0.3596	2.256	969.9917	0.2262	2.035	973.0006	0.4100	2.215
969.9707	0.1901	2.216	972.9762	0.3681	2.276	969.9975	0.2361	2.024	973.0053	0.4183	2.225
969.9790	0.2044	2.200	972.9805	0.3755	2.288	970.0032	0.2460	2.027	973.0108	0.4276	2.231
969.9846	0.2139	2.204	972.9850	0.3832	2.293	970.0143	0.2651	2.024	973.0204	0.4442	2.266
969.9907 969.9965	0.2245	2.191 2.188	972.9897 972.9955	0.3913	2.303	970.0217 970.0284	0.2778	2.026 2.046	973.0257 973.0314	0.4533 0.4632	2.276
970.0022	0.2443	2.173	972.9998	0.4013	2.331	770.0284	0.2074	2.046	973.0362	0.4715	
970.0133	0.2635	2.181	973.0046	0.4169	2.370	970.8690	0.7376	2.031	973.0422		2.337
970.0207	0.2762	2.185	973.0101	0.4264	2.392	970.8752	0.7483	2.019	973.0477	0.4912	
970.0274	0.2877	2.182	973.0197	0.4429	2.420	970.9349	0.8513	2.106			
			973.0249	0.4520	2.442	970.9425	0.8643	2.133			
970.8680	0.7359	2.176	973.0307	0.4619	2.449	970.9478	0.8735	2.149	НJD		
970.8742	0.7466	2.174	973.0355	0.4702	2.463	970.9532	0.8827	2.163	2445000+	Phase	ΔU
970.9339	0.8496	2.244	973.0412 973.0468	0.4801 0.4896	2.495	970.9591 970.9644	0.8929	2.202			
970.9402 970.9415	0.8604 0.8626	2.273 2.275	713.0400	J. 4070	2.451	970.9696	0.9020	2.237 2.253			
970.9413	0.8718	2.296				970.9753	0.9208	2.306	958.8454	0.0228	2.819
970.9522	0.8810	2.317				970.9810	0.9307	2.377	959.0334	0.3466	2.304
970.9581	0.8912	2.348	HJD	Phase	ΔΒ	970.9875	0.9418	2.444			
970.9634	0.9003	2.374	2445000+			970.9931	0.9514	2.517	971.7332	0.2266	2.054
970.9687	0.9094	2.388	-			970.9989	0.9614	2.608	971.7501	0.2557	2.052
970.9743	0.9191	2.445	050 0445	0 0007	2 722	971.0044	0.9709	2.670	072 0005	0 05:5	
970.9800	0.9289	2.508	958.8442	0.0207	2.782	971.0118 971.0176	0.9836	2.768	972.8989 972.9059	0.2349	2.064
970.9865 970.9921	0.9401	2.564	958.8638 958.8730	0.0545	2.557 2.428	971.0178	0.9937 0.9958	2.793 2.810	972.9039	0.2469 0.2581	2.063
970.9921	0.9599	2.698	958.8815		2.335	971.0235		2.807	116.7164	0.2301	2.066
	20.2//							·			